

EVOLVING SURFACE FINITE ELEMENT METHODS FOR RANDOM ADVECTION-DIFFUSION EQUATIONS

ANA DJURDJEVAC, CHARLES M. ELLIOTT, RALF KORNUBER,
AND THOMAS RANNER

ABSTRACT. Even though random partial differential equations (PDEs) have become a very active field of research, the analysis and numerical analysis of random PDEs on surfaces still appears to be in its infancy (see, however [16]). In this paper, we introduce and analyse a surface finite element discretization of advection-diffusion equations with uncertain coefficients on evolving hypersurfaces. After stating unique solvability of the resulting semi-discrete problem, we prove optimal error bounds for the semi-discrete solution and Monte Carlo samplings of its expectation in appropriate Bochner spaces. Our theoretical findings are illustrated by numerical experiments in two and three space dimensions.

1. INTRODUCTION

During the last decades surface partial differential equations (PDEs), i.e. PDEs on stationary or evolving curved domains have grown in importance. The numerical analysis of surface PDEs started with the paper [17] and later it developed in [19], [15], [39], etc. Dziuk and Elliott have introduced the evolving surface finite element method for PDEs on moving hypersurfaces ([18], [20]). Recently this topic has been generalized ([1], [2]) to a more abstract level, i.e. to parabolic PDEs posed in an evolving Hilbert space.

Surface PDEs appear in a variety of applications such as image processing [29], computer graphics [7], biological modelling [31] and engineering [35]. Particularly for this paper, the motivating example is modelling the transport of a surface active agent (surfactant) on the interface between two fluids ([38], [28]).

In all these applications uncertainty naturally appears, for example when the input data is not precisely known, either because direct measurement are not practical or they admit intrinsic variability. Therefore, even when a model is deterministic, it is not enough to consider just one simulation since the parameters that are appearing are often estimated from measurements and are subject to uncertainty. A probabilistic approach to deal with this kind of uncertainty is to present input data as a random field over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. This approach has been widely used in the field uncertainty quantification, especially considering random PDEs i.e. PDEs with random coefficients (the general reference here are [34] and [32]). In this work we address specifically parabolic random PDEs, which have been so far studied in several papers ([36], [27], [9], [30]).

2010 *Mathematics Subject Classification.* 65N12, 65N30, 65C05.

The research of TR was funded by the Engineering and Physical Sciences Research Council (EPSRC EP/J004057/1).

This motivates us to consider random surface PDEs. The first analysis, up to our knowledge, of the random PDEs on moving hypersurfaces is given in [16]. However the numerical analysis was not considered, and it will be the focus of this paper. More specifically, we will consider the numerical analysis of a semi-discretization of

$$\partial^\bullet u - \nabla_\Gamma(\alpha \nabla_\Gamma u) + u \nabla_\Gamma \cdot \mathbf{v} = 0$$

on evolving compact hypersurface $\Gamma(t) \subset \mathbb{R}^n, n = 1, 2$ for time $t \in [0, T], T > 0$ with given uniformly bounded random coefficient α and deterministic velocity $\mathbf{v}$. Therefore, our solution u is also a random variable. Here ∇_Γ is a tangential gradient and $\partial^\bullet$ is a material derivative that is defined path-wise in a standard way. This is an advection-diffusion equation that arises from conservation of a random variable u on an evolving hypersurface. Its path-wise derivation can be found for example in [18].

Since we assume the uniform bounds for the random diffusion coefficient α , we will consider mean-weak formulation of (1)

$$(1) \quad \int_{\Omega} \int_{\Gamma(t)} \partial^\bullet u(t) \varphi + \int_{\Omega} \int_{\Gamma(t)} \alpha(t) \nabla_\Gamma u(t) \cdot \nabla_\Gamma \varphi + \int_{\Omega} \int_{\Gamma(t)} u(t) \varphi \nabla_\Gamma \cdot \mathbf{v}(t) = 0$$

where φ is an arbitrary random test function defined on the surface $\Gamma(t)$ for every t and has bounded second moment.

The well-posedness of the problem and regularity results have been shown in [16]. Here we will focus on its discretization. For discretization in space we will use evolving surface finite element method (ESFEM) that was proposed in [18] using piecewise linear finite element functions on a triangulated surface interpolating $\Gamma(t)$ whose vertices move with velocity $\mathbf{v}$. An optimal L^2 -estimate for the error between continuous and semi-discrete problem in the deterministic case was proved in [20]. We will use similar analysis for discretization in space of (1). The challenges for convergence analysis lie in proving the well-posedness in semi-discrete case and controlling the constants that appear in error bounds and showing its independence on a path. The main difficulty is to define the path-wise Ritz projection and prove its integrability with respect to probability $\mathbb{P}$ and boundedness of its second moments, which will lead to an error bound for projection.

In order to deal with uncertainty in our numerical scheme we will use the standard Monte Carlo (MC) method. It has dimensional independent convergence rate of the sample error. This is especially useful for high-dimensional problems i.e. for problems that depend on a large number of independent random variables, and this case often can not be well treated with other methods. However, MC convergence rate is very slow. Namely it behaves as $M^{-1/2}$, where M is the total number of realizations. Therefore many other techniques have been developed in order to improve the convergence rate, such as variance reduction techniques. One of those widely used methods is Multilevel MC, applied especially to random PDEs in [3], [10], [12] etc. Nevertheless, since there is no other available numerical analysis of the equation (1) and because of the simplicity of implementation of MC method, we will consider just MC as a starting point and consider possible improvements in the future research.

The paper is organized as follows. We start the second section by setting up the notation, description of the hypersurfaces, defining functional spaces and material derivative and formulating the problem. It mainly contains the framework defined

in [16] that we will use. In the third section we proceed with introducing the ES-FEM following [18]. Section 4 contains the precise formulation of the semi-discrete problem and its well-posedness. In the fifth section we give results concerning the approximation of the surface. In particular we show that estimates from deterministic setting will still hold with constants independent of the realization. We define and prove regularity of path-wise Ritz projection. The section ends with the L^2 -error bound for the ESFEM. In Section 6 we prove the main result on error bound of MC-ESFEM. In the final section we illustrate our results by numerical experiments.

2. RANDOM ADVECTION-DIFFUSION EQUATIONS ON EVOLVING HYPERSURFACES

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space with sample space Ω , a σ -algebra of events $\mathcal{F}$ and a probability $\mathbb{P}: \mathcal{F} \rightarrow [0, 1]$. In addition, we assume that $L^2(\Omega)$ is a separable space. For this assumption it suffices to assume that $(\Omega, \mathcal{F}, \mathbb{P})$ is separable [25, Exercise 43.(1)]. We will only consider a fixed finite time interval $[0, T]$, where $T \in (0, \infty)$. Furthermore, we will denote by $\mathcal{D}([0, T]; V)$ the space of infinitely differentiable functions with values in a Hilbert space V and compact support in $(0, T)$.

2.1. Hypersurfaces. We first recall some basic notions and results concerning hypersurfaces and Sobolev spaces on hypersurfaces. We refer to [13] and [22] for more details.

Let $\Gamma \subset \mathbb{R}^{n+1}$ ($n = 1, 2$) be a $\mathcal{C}^2$ compact, connected, orientable, n -dimensional hypersurface. For a function $f: \Gamma \rightarrow \mathbb{R}$ allowing for a differentiable extension $\tilde{f}$ to an open neighbourhood of Γ in $\mathbb{R}^{n+1}$ we define the *tangential gradient* by

$$(2) \quad \nabla_{\Gamma} f(x) := \nabla \tilde{f}(x) - \nabla \tilde{f}(x) \cdot \nu(x) \nu(x), \quad x \in \Gamma,$$

where $\nu(x)$ denotes the unit normal to Γ . Note that $\nabla_{\Gamma} f(x)$ is the orthogonal projection of $\nabla \tilde{f}$ onto the tangent space to Γ at x (thus a tangential vector). It depends only on the values of $\tilde{f}$ on Γ [22, Lemma 2.4], which makes the definition (2) independent of the extension $\tilde{f}$. The tangential gradient is a vector-valued quantity and for its components we will use the notation $\nabla_{\Gamma} f(x) = (\underline{D}_1 f(x), \dots, \underline{D}_{n+1} f(x))$. Now we can define the *Laplace-Beltrami* operator by

$$\Delta_{\Gamma} f(x) = \nabla_{\Gamma} \cdot \nabla_{\Gamma} f(x) = \sum_{i=1}^{n+1} \underline{D}_i \underline{D}_i f(x), \quad x \in \Gamma.$$

We will consider weak formulations of PDEs on Γ , which involve the concept of Sobolev spaces on surfaces. We define $L^2(\Gamma)$ as a set of all measurable functions $f: \Gamma \rightarrow \mathbb{R}$ such that $\|f\|_{L^2(\Gamma)} := (\int_{\Gamma} |f(x)|^2)^{1/2}$ is finite. We say that a function $f \in L^2(\Gamma)$ has a weak derivative $g_i = D_i f \in L^2(\Gamma)$, ($i = \{1, \dots, n+1\}$), if for every function $\phi \in \mathcal{C}^1(\Gamma)$ and every i there holds

$$\int_{\Gamma} f \underline{D}_i \phi = - \int_{\Gamma} \phi g_i + \int_{\Gamma} f \phi H \nu_i.$$

The Sobolev space on Γ is defined by

$$H^1(\Gamma) = \{f \in L^2(\Gamma) \mid D_i f \in L^2(\Gamma), i = 1, \dots, n+1\}$$

with the norm $\|f\|_{H^1(\Gamma)} = \sqrt{\|f\|_{L^2(\Gamma)}^2 + \|\nabla_{\Gamma} f\|_{L^2(\Gamma)}^2}$.

For a description of evolving hypersurfaces we will consider two approaches. We first introduce evolutions according to a given velocity field v . To this end, we assume that for every $t \in [0, T]$, $\Gamma(t)$ satisfies the same properties as Γ and we set $\Gamma_0 := \Gamma(0)$. Furthermore, we assume the existence of a flow

$$\Phi_t^0(\cdot) := \Phi(\cdot, t): \Gamma_0 \rightarrow \Gamma(t), \quad \Phi \in \mathcal{C}^1([0, T], \mathcal{C}^1(\Gamma_0)) \cap \mathcal{C}^0([0, T], \mathcal{C}^3(\Gamma_0))$$

which is a diffeomorphism that satisfies

$$\frac{d}{dt} \Phi_t^0(\cdot) = v(t, \Phi_t^0(\cdot)), \quad \Phi_0^0(\cdot) = \text{Id}(\cdot),$$

where $v: [0, T] \times \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ is a $\mathcal{C}^2$ -velocity field with uniformly bounded divergence

$$(3) \quad |\nabla_{\Gamma(t)} \cdot v(t)| \leq C \quad \text{for all } t \in [0, T].$$

For the numerical analysis it is sometimes convenient to represent $\Gamma(t)$ as a zero level set of a suitable function defined on a subset of $\mathbb{R}^{n+1}$. Under the given regularity assumptions for $\Gamma(t)$, it follows by Jordan-Brouwer theorem that $\Gamma(t)$ is the boundary of an open bounded domain. Thus, $\Gamma(t)$ can be represented as the zero level set

$$\Gamma(t) = \{x \in \mathcal{N}(t) \mid d(x, t) = 0\}, \quad t \in [0, T],$$

of a signed distance function $d = d(x, t)$ defined on an open neighborhood $\mathcal{N}(t)$ of $\Gamma(t)$ such that $|\nabla d| \neq 0$ and $t \in [0, T]$. Note that $d, d_t, d_{x_i}, d_{x_i x_j} \in \mathcal{C}^1(\mathcal{N}_T)$, $i, j = 1, \dots, n+1$ holds on $\mathcal{N}_T = \bigcup_t \mathcal{N}(t) \times \{t\}$. We also choose $\mathcal{N}(t)$ such that for every $x \in \mathcal{N}(t)$ and $t \in [0, T]$ there exists unique $p(x, t) \in \Gamma(t)$ such that

$$(4) \quad x = p(x, t) + d(x, t)\nu(p(x, t), t),$$

and fix the orientation of $\Gamma(t)$ by choosing the normal vector field $\nu(x, t) := \nabla d(x, t)$. We will use (4) below to define the lift of functions on approximate hypersurfaces.

2.2. Function spaces. In order to treat the evolving spaces, we need to define special Bochner-type function spaces of random functions that are defined on evolving spaces for which we have that $u(\omega, t): \Gamma(t) \rightarrow \mathbb{R}$. The definition of these spaces uses the standard idea to map each domain at time t to the fixed initial domain Γ_0 using the flow Φ_t^0 . This approach is similar to Arbitrary Lagrangian Eulerian (ALE) framework. The results of this and next subsection can be found in [1] that gives the general framework for parabolic PDEs on evolving spaces and [16] where the probability setting is given.

For each $t \in [0, T]$, let us define

$$(5) \quad V(t) := L^2(\Omega, H^1(\Gamma(t))) \cong L^2(\Omega) \otimes H^1(\Gamma(t))$$

$$(6) \quad H(t) := L^2(\Omega, L^2(\Gamma(t))) \cong L^2(\Omega) \otimes L^2(\Gamma(t))$$

where the isomorphism hold because all considered spaces are separable Hilbert spaces (see [37]). The dual space of $V(t)$ is the space $V^*(t) = L^2(\Omega, H^{-1}(\Gamma(t)))$, where $H^{-1}(\Gamma(t))$ is the dual space of $H^1(\Gamma(t))$. Using the tensor product structure of these spaces [24, Lemma 4.34], it follows that $V(t) \subset H(t) \subset V^*(t)$ is a Gelfand triple for every $t \in [0, T]$.

Remark 1. For convenience we will often (but not always) write $u(\omega, x)$ instead of $u(\omega)(x)$, which is justified by the tensor structure of the spaces.

For an evolving family of Hilbert spaces $X = (X(t))_{t \in [0, T]}$, we connect the space $X(t)$ at any time $t \in [0, T]$ with fixed initial space $X(0)$, using the family of maps $\phi_t : X(0) \rightarrow X(t)$, which we call the pushforward map. More precisely we use its inverse map $\phi_{-t} : X(t) \rightarrow X(0)$ that we call the pullback map. We define the pullback operator $\phi_{-t} : L^2(\Omega, L^2(\Gamma(t))) \rightarrow L^2(\Omega, L^2(\Gamma_0))$ in the natural way, using the parametrisation Φ_t^0 that maps Γ_0 onto $\Gamma(t)$

$$(\phi_{-t}u)(\omega, x) := u(\omega, \Phi_t^0(x)) \quad \text{for every } x \in \Gamma(0), \omega \in \Omega.$$

We define general Bochner-type spaces of functions that are defined on evolving spaces as in [1]:

$$L_X^2 := \left\{ u : [0, T] \rightarrow \bigcup_{t \in [0, T]} X(t) \times \{t\}, t \mapsto (\bar{u}(t), t) \mid \phi_{-(\cdot)} \bar{u}(\cdot) \in L^2(0, T; X_0) \right\}$$

$$L_{X^*}^2 := \left\{ f : [0, T] \rightarrow \bigcup_{t \in [0, T]} X^*(t) \times \{t\}, t \mapsto (\bar{f}(t), t) \mid \phi_{(\cdot)}^* \bar{f}(\cdot) \in L^2(0, T; X_0^*) \right\}.$$

Remark 2. In the following we will identify $u(t) = (\bar{u}(t); t)$ with $\bar{u}(t)$.

From [1, Lemma 2.15] it follows that $L_{X^*}^2$ and $(L_X^2)^*$ are isometrically isomorphic.

The spaces L_X^2 and $L_{X^*}^2$ are separable Hilbert spaces [1, Corollary 2.11] with the inner product defined as

$$(u, v)_{L_X^2} = \int_0^T (u(t), v(t))_{X(t)} dt \quad (f, g)_{L_{X^*}^2} = \int_0^T (f(t), g(t))_{X^*(t)} dt.$$

For the evolving families V and H defined by (5) and (6), from [16, Lemma 3.5] it follows that the spaces L_V^2 , $L_{V^*}^2$ and L_H^2 are well-defined and

$$L_V^2 \subset L_H^2 \subset L_{V^*}^2$$

is a Gelfand triple that we will consider for our equation.

2.3. Material derivative and solution space. In this subsection we define a material derivative for random functions that also takes into account the spatial movement in the same way as in [16]. Since we focus on numerical analysis in this paper, we consider sufficiently smooth solution and it is enough to define a (strong) material derivative for continuous differentiable functions. First let us define the spaces of pushed-forward continuously differentiable functions

$$\mathcal{C}_V^j := \{u \in L_V^2 \mid \phi_{-(\cdot)} u(\cdot) \in \mathcal{C}^j([0, T], L^2(\Omega, H^1(\Gamma_0)))\} \quad \text{for } j \in \{0, 1, 2\}.$$

For $u \in \mathcal{C}_V^1$ the material derivative $\partial^\bullet u \in \mathcal{C}_V^0$ is defined by

$$(7) \quad \partial^\bullet u := \phi_t \left(\frac{d}{dt} \phi_{-t} u \right) = u_t + \nabla u \cdot v.$$

It would be more precise to define the material derivative of u using its smooth extension $\tilde{u}$ in $\mathcal{N}_T$ for which $\nabla \tilde{u}$ and $\tilde{u}_t$ are well defined. Since it can be shown that because of the smoothness of $\Gamma(t)$ and evolution Φ_t^0 this definition is independent of the choice of extension $\tilde{u}$, we abuse notation and use the same u .

Remark 3. It is possible to define a weak material derivative that needs less regularity [16, Definition 3.9].

Now we define the solution space as

$$(8) \quad W(V, H) := \{u \in L_V^2 \mid \partial^\bullet u \in L_H^2\}.$$

The solution space $W(V, H)$ is a Hilbert space with the inner product defined via

$$(u, v)_{W(V, H)} = \int_0^T \int_\Omega (u, v)_{H^1(\Gamma(t))} + \int_0^T \int_\Omega (\partial^\bullet u, \partial^\bullet v)_{L^2(\Gamma(t))}.$$

2.4. Problem formulation and well-posedness. We consider an initial value problem for an advection-diffusion equation on the evolving surface $\Gamma(t)$, $t \in [0, T]$, which in strong form reads

$$(9) \quad \begin{aligned} \partial^\bullet u - \nabla_\Gamma \cdot (\alpha \nabla_\Gamma u) + u \nabla_\Gamma \cdot \mathbf{v} &= f \\ u(0) &= u_0 \end{aligned}$$

Here the diffusion coefficient α and the initial function u_0 are random functions, and we set $f \equiv 0$ for ease of presentation.

A *weak solution* of (9) amounts to find $u \in W(V, H)$ that satisfies the initial condition $u(0) = u_0$ and

$$(10) \quad \int_\Omega \int_{\Gamma(t)} \partial^\bullet u(t) \varphi + \int_\Omega \int_{\Gamma(t)} \alpha(t) \nabla_\Gamma u(t) \cdot \nabla_\Gamma \varphi + \int_\Omega \int_{\Gamma(t)} u(t) \varphi \nabla_\Gamma \cdot \mathbf{v}(t) = 0,$$

for every $\varphi \in L^2(\Omega, H^1(\Gamma(t)))$ and a.e. $t \in [0, T]$.

Existence and uniqueness will follow from the following assumptions.

Assumption 1. *The diffusion coefficient α satisfies the following conditions*

- a) $\alpha: \Omega \times \mathcal{G}_T \rightarrow \mathbb{R}$ is a $\mathcal{F} \otimes \mathcal{B}(\mathcal{G}_T)$ -measurable, where $\mathcal{G}_T := \bigcup_t \Gamma(t) \times \{t\}$ is a space-time domain.
- b) $\alpha(\omega, \cdot, \cdot) \in C^1(\mathcal{G}_T)$ holds for $\mathbb{P}$ -a.e. $\omega \in \Omega$, which implies boundedness of $|\partial^\bullet \alpha(\omega)|$ on $\mathcal{G}_T$ and in addition we assume that this bound is uniform in ω .
- c) α is uniformly bounded from above and below in the sense that there exist positive constants $\alpha_{\min}$ and $\alpha_{\max}$ such that

$$(11) \quad 0 < \alpha_{\min} \leq \alpha(\omega, x, t) \leq \alpha_{\max} < \infty \quad \forall (x, t) \in \mathcal{G}_T$$

holds for $\mathbb{P}$ -a.e. $\omega \in \Omega$.

and the initial function satisfies $u_0 \in L^2(\Omega, L^2(\Gamma_0))$.

We will need the following assumption about the uniform bound for the tangential gradient of the diffusion coefficient in order to get the $L^2(\Gamma(t))$ bound.

Assumption 2. *Assume that there exists a constant C independent of ω such that*

$$|\nabla_\Gamma \alpha(\omega, x, t)| \leq C \quad \text{for all } (x, t) \in \mathcal{G}_T$$

holds for $\mathbb{P}$ -a.e. $\omega \in \Omega$.

Note that from (11) and Assumption 2 it follows that $\|\alpha(\omega, t)\|_{C^1(\bar{\Gamma}(t))}$ is uniformly bounded i.e. independently of ω , which we will use to obtain $H^2(\Gamma(t))$ -bound.

It follows from [16, Theorem 4.9] that Assumption 1 provides existence of a unique solution $u \in W(V, H)$ of (10) satisfying the *a priori* bound

$$\|u\|_{W(V, H)} \leq C \|u_0\|_{V_0},$$

with a constant $C \in \mathbb{R}$. In the error analysis, we will assume further that u has a path-wise strong material derivative. This is for all $\omega \in \Omega$, $u(\omega) \in C_V^1$.

In order to derive a more convenient formulation of (10), we introduce the bilinear forms

$$(12) \quad \begin{aligned} m(u, \varphi) &:= \int_{\Omega} \int_{\Gamma(t)} u \varphi, & g(v; u, \varphi) &:= \int_{\Omega} \int_{\Gamma(t)} u \varphi \nabla_{\Gamma} \cdot v, \\ a(u, \varphi) &:= \int_{\Omega} \int_{\Gamma(t)} \alpha \nabla_{\Gamma} u \cdot \nabla_{\Gamma} \varphi & b(v; u, \varphi) &= \int_{\Omega} \int_{\Gamma(t)} B(\omega, v) \nabla_{\Gamma} u \cdot \nabla_{\Gamma} \varphi. \end{aligned}$$

for $u, \varphi \in L^2(\Omega, H^1(\Gamma(t)))$ and each $t \in [0, T]$. The tensor B in the definition of $b(v; u, \varphi)$ takes the form $B(\omega, v) = (\partial^{\bullet} \alpha + \alpha \nabla_{\Gamma} \cdot v) \mathbb{I} - 2\alpha D_{\Gamma}(v)$ denoting $(D_{\Gamma} v)_{ij} = \underline{D}_j v^i$. Note that (3) and the uniform boundedness of $\partial^{\bullet} \alpha$ on $\mathcal{G}_T$ imply that $|B(\omega, v)| \leq C$ holds $\mathbb{P}$ -a.e. $\omega \in \Omega$ with some $C \in \mathbb{R}$.

The transport formulae for the differentiation of time dependent surface integral then read (see e.g. [16])

$$(13) \quad \frac{d}{dt} m(u, \varphi) = m(\partial^{\bullet} u, \varphi) + m(u, \partial^{\bullet} \varphi) + g(v; u, \varphi),$$

$$(14) \quad \frac{d}{dt} a(u, \varphi) = a(\partial^{\bullet} u, \varphi) + a(u, \partial^{\bullet} \varphi) + b(v; u, \varphi)$$

where the first relation holds in $W(H, H)$ and the second in $W(V, V)$.

Using (13), we can rewrite (10) briefly as

$$(15) \quad \frac{d}{dt} m(u, \varphi) + a(u, \varphi) = m(u, \partial^{\bullet} \varphi) \quad \text{for all } \varphi \in W(V, H).$$

This formulation will be used in the sequel.

3. EVOLVING SURFACE FINITE ELEMENT APPROXIMATION

In this section we introduce the notation for the evolving surface finite element method (ESFEM) following [18].

3.1. Evolving triangulated surfaces. Let $\Gamma_{h,0}$ be a polyhedral approximation of Γ_0 that lies in $\mathcal{N}(0)$ with nodes $\{X_{j,0}\}_{j=1}^N$ on Γ_0 . We evolve the nodes by the smooth surface velocity $\dot{X}_j(t) = v(X_j(t), t)$ and $X_j(0) = X_j^0$, $j = 1, \dots, N$. For each $t \in [0, T]$, linear interpolation between these nodes defines a family of discrete surfaces $\{\Gamma_h(t)\}$. We denote by $\mathcal{T}_h(t)$ a triangulation of $\Gamma_h(t)$ consisting of closed simplices and assume that it is quasi-uniform uniformly in time. We define h as the maximal diameter of elements in $\mathcal{T}_h(t)$ uniformly in time $h := \sup_{t \in (0, T)} \max_{E(t) \in \mathcal{T}_h(t)} \text{diam } E(t)$. Previous assumptions ensure that $\Gamma_h(t)$ can be considered as interpolation of $\Gamma(t)$ and by relation (4) there is a bijective correspondence between the triangles on $\Gamma_h(t)$ and the induced curve linear triangles on $\Gamma(t)$. We will denote by $\mathcal{G}_T^h$ the discrete space time domain given by

$$\mathcal{G}_T^h := \bigcup_t \Gamma_h(t) \times \{t\}.$$

We define the gradient of $\eta_h: \Gamma_h(t) \rightarrow \mathbb{R}$ in an element-wise sense

$$\nabla_{\Gamma_h} \eta_h|_E = \nabla \eta_h - \nabla \eta_h \cdot \nu_h \nu_h, \quad \text{for all } E \in \mathcal{T}_h(t)$$

where ν_h is the unit outward pointing normal to E .

For continuous functions $\eta_h(\cdot, t): \Gamma_h(t) \rightarrow \mathbb{R}$, we define its lift $\eta_h^l(\cdot, t): \Gamma(t) \rightarrow \mathbb{R}$ onto $\Gamma(t)$ using (4) by

$$\eta_h^l(p, t) := \eta_h(x(p, t), t), \quad p \in \Gamma(t).$$

Any continuous function $\eta: \Gamma(t) \rightarrow \mathbb{R}$ can be extended to $\mathcal{N}(t)$ in the normal direction by $\eta^{-l}(x, t) := \eta(p(x, t), t)$. Therefore, for η we define its inverse lift $\eta^{-l}: \Gamma_h(t) \rightarrow \mathbb{R}$ by

$$\eta^{-l}(x, t) := \eta(x - d(x, t)\nu(x, t), t), \quad x \in \Gamma_h(t).$$

It follows that $(\eta^{-l})^l = \eta$.

The next lemma gives the relation between norms on continuous and discrete hypersurface. The analogue results hold in the deterministic case, see for [17] for the proof.

Lemma 1. *For $t \in [0, T]$, $\omega \in \Omega$, let $\eta_h(\omega): \Gamma_h \rightarrow \mathbb{R}$ and denote its lift by $\eta_h^l(\omega): \Gamma \rightarrow \mathbb{R}$. Then for corresponding plane simplex $E \subset \Gamma_h(t)$ and curvilinear simplex $e \subset \Gamma(t)$, there is a constant $c > 0$ independent of h , t and ω s.t.*

$$(16) \quad \frac{1}{c} \|\eta_h\|_{L^2(\Omega, L^2(E))} \leq \|\eta_h^l\|_{L^2(\Omega, L^2(e))} \leq c \|\eta_h\|_{L^2(\Omega, L^2(E))}$$

$$(17) \quad \frac{1}{c} \|\nabla_{\Gamma_h} \eta_h\|_{L^2(\Omega, L^2(E))} \leq \|\nabla_{\Gamma} \eta_h^l\|_{L^2(\Omega, L^2(e))} \leq c \|\nabla_{\Gamma_h} \eta_h\|_{L^2(\Omega, L^2(E))}$$

$$(18) \quad \frac{1}{c} \|\nabla_{\Gamma_h}^2 \eta_h\|_{L^2(\Omega, L^2(E))} \leq c \|\nabla_{\Gamma}^2 \eta_h^l\|_{L^2(\Omega, L^2(e))} + ch \|\nabla_{\Gamma} \eta_h^l\|_{L^2(\Omega, L^2(e))},$$

if the norms exist.

3.2. Finite element space and discrete velocities. For every $t \in [0, T]$ we define a *finite element space* and its lift by

$$(19) \quad S_h(t) := \{\phi \in \mathcal{C}^0(\Gamma_h(t)) \mid \phi|_e \text{ is linear affine for each } e \in \mathcal{T}_h(t)\}$$

$$(20) \quad S_h^l(t) := \{\phi^l \in \mathcal{C}^0(\Gamma(t)) \mid \phi^l = \phi(x(a)), \phi \in S_h(t)\}.$$

Note that $S_h(t) \subset H^1(\Gamma_h(t))$ and that there is a unique correspondence between each element from $S_h(t)$ and $S_h^l(t)$. We denote by $\{\chi_j(t)\}_{j=1, \dots, N}$ the nodal basis of $S_h(t)$, such that $\chi_j(X_i(t), t) = \delta_{ij}$.

Definition 1. *We define a discrete material velocity V_h for $x = X(t)$ on the surface $\Gamma_h(t)$ by*

$$X'(t) = V_h(X(t), t) = I_h v(X(t), t) := \sum_{j=1}^N X_j'(t) \chi_j^t(X(t)).$$

Then an associated discrete material velocity v_h for $Y(t) = a(X(t), t)$ on $\Gamma(t)$ is defined by

$$Y'(t) = v_h(Y(t), t) = \frac{\partial a}{\partial t}(X(t), t) + V_h(X(t), t) \cdot \nabla a(X(t), t).$$

3.3. Defining problem on polyhedral surfaces. After introducing all the necessary definitions for the evolving triangulated family, we can define the problem on $\Gamma_h(t)$ and state the analogue well-posedness theorem and transport theorem for $\Gamma_h(t)$. The corresponding Gelfand triple on $\Gamma_h(t)$ is naturally

$$L^2(\Omega, H^1(\Gamma_h(t))) \subset L^2(\Omega, L^2(\Gamma_h(t))) \subset L^2(\Omega, H^{-1}(\Gamma_h(t))).$$

We will denote

$$\mathcal{V}_h(t) := L^2(\Omega, H^1(\Gamma_h(t))) \quad \text{and} \quad \mathcal{H}_h(t) := L^2(\Omega, L^2(\Gamma_h(t))).$$

Similarly as in the continuous case, we derive the Gelfand triple of evolving Bochner-Sobolev spaces

$$L^2_{\mathcal{V}_h(t)} \subset L^2_{\mathcal{H}_h(t)} \subset L^2_{\mathcal{V}_h^*(t)}.$$

Analogy to definition (7), we define the discrete material derivatives for smooth enough functions $\phi_h \in L^2_{\mathcal{V}_h}$ and $\varphi_h \in L^2_V$ element by element

$$\partial_h^\bullet \phi_h|_{E(t)} := (\phi_{ht} + V_h \cdot \nabla \phi_h)|_{E(t)} \quad \text{and} \quad \partial_h^\bullet \varphi_h|_{e(t)} := (\varphi_{ht} + v_h \cdot \nabla \varphi_h)|_{e(t)},$$

for any $E(t) \subset \Gamma_h(t)$ and $e(t) = p(E(t), t) \subset \Gamma(t)$.

Remark 4. *The basis functions of the evolving finite element space satisfy the transport property [20, Lemma 4.1]*

$$(21) \quad \partial_h^\bullet \chi_j = 0 \quad \partial_h^\bullet \chi_j^l = 0.$$

A solution space on $\{\Gamma_h(t)\}_{t \in [0, T]}$ is defined in an analogue to (8) by

$$W(\mathcal{V}_h, \mathcal{H}_h) = \{u \in L^2_{\mathcal{V}_h} \mid \partial_h^\bullet u \in L^2_{\mathcal{H}_h}\}.$$

We define the discrete analogues to the bilinear forms introduced in (12) on $\mathcal{V}_h(t) \times \mathcal{V}_h(t)$

$$\begin{aligned} m_h(u_h, \varphi_h) &:= \int_{\Omega} \int_{\Gamma_h(t)} u_h \varphi_h, & a_h(u_h, \varphi_h) &:= \int_{\Omega} \int_{\Gamma_h(t)} \alpha^{-l} \nabla_{\Gamma_h} u_h \cdot \nabla_{\Gamma_h} \varphi_h, \\ g_h(V_h; u_h, \varphi_h) &:= \int_{\Omega} \int_{\Gamma_h(t)} u_h \varphi_h \nabla_{\Gamma_h} \cdot V_h, \end{aligned}$$

Here, we denote by α^{-l} the inverse lift of α , which satisfies a discrete version of Assumption 1. In particular, α^{-l} is an $\mathcal{F} \otimes \mathcal{B}(\mathcal{G}_T^h)$ -measurable function, $\alpha^{-l}(\omega, \cdot, \cdot)|_{E_T} \in \mathcal{C}^1(E_T)$ for all space-time elements $E_T := \bigcup_t E(t) \times \{t\}$, and $\alpha_{\min} \leq \alpha^{-l}(\omega, x, t) \leq \alpha_{\max}$ for all $\omega \in \Omega$, $(x, t) \in \mathcal{G}_T^h$.

Since the probability space does not depend on time, the discrete analogue of the transport formulae hold, where the discrete material velocity and discrete tangential gradients appear and are understood in an element-wise sense. The discrete result is stated for example in [20, Lemma 4.2], in addition one should integrate over Ω . We state the result for completeness.

Lemma 2 (Transport theorems for triangulated surfaces). *Let $\Gamma_h(t)$ be an evolving admissible triangulation with material velocity V_h . Then*

$$\frac{d}{dt} \int_{\Omega} \int_{\Gamma_h(t)} f = \int_{\Omega} \int_{\Gamma_h(t)} \partial_h^\bullet f + f \nabla_{\Gamma_h} \cdot V_h.$$

For $\phi, U_h \in \mathcal{V}_h(t)$,

$$\begin{aligned} \frac{d}{dt} m_h(\phi, U_h) &= m(\partial_h^\bullet \phi, U_h) + m(\phi, \partial_h^\bullet U_h) + g_h(V_h; \phi, U_h) \\ \frac{d}{dt} a_h(\phi, U_h) &= a(\partial_h^\bullet \phi, U_h) + a(\phi, \partial_h^\bullet U_h) + b(V_h; \phi, U_h) \end{aligned}$$

with the bilinear form

$$b(V_h; \phi, U_h) = \sum_{E(t) \in \mathcal{T}_h(t)} \int_{\Omega} \int_{E(t)} B_h(\omega, V_h) \nabla_{\Gamma_h} \phi \cdot \nabla_{\Gamma_h} U_h$$

and the tensor

$$B_h(\omega, V_h) = \partial_h^\bullet \alpha^{-l} + \alpha^{-l} \nabla_{\Gamma_h} \cdot V_h - 2\alpha^{-l} D_h(V_h)$$

where $(D_h(V_h))_{ij} = \underline{D}_{h,j} V_h^i$.

Let $\Gamma(t)$ be an evolving surface decomposed into curved elements whose edges move with velocity v_h . Then

$$\frac{d}{dt} \int_{\Omega} \int_{\Gamma(t)} f = \int_{\Omega} \int_{\Gamma(t)} \partial_h^\bullet f + f \nabla_\Gamma \cdot v_h.$$

For $\varphi, u_h, \partial_h^\bullet \varphi, \partial_h^\bullet u_h \in V(t)$,

$$\begin{aligned} \frac{d}{dt} m(\varphi, u_h) &= m(\partial_h^\bullet \varphi, u_h) + m(\varphi, \partial_h^\bullet u_h) + g(v_h; \varphi, u_h) \\ \frac{d}{dt} a(\varphi, u_h) &= a(\partial_h^\bullet \varphi, u_h) + a(\varphi, \partial_h^\bullet u_h) + b(v_h; \varphi, u_h). \end{aligned}$$

The next lemma gives us the uniform bound for the divergence of V_h and tensor B_h (see [23, Lemma 3.3.1]).

Lemma 3. *Under previous assumptions on $\{\Gamma_h(t)\}_t$, it holds*

$$\sup_{t \in [0, T]} (\|\nabla_{\Gamma_h} \cdot V_h\|_{L^\infty(\Gamma_h(t))} + \|B_h\|_{L^2(\Omega, L^\infty(\Gamma_h(t)))}) \leq c \sup_{t \in [0, T]} \|v\|_{C^2(\mathcal{N}_T)}.$$

The existence and uniqueness of the weak solution $U_h \in W_h(\mathcal{V}_h, \mathcal{H}_h)$ of the equation on the polyhedral surface

$$(22) \quad \frac{d}{dt} m_h(U_h, \phi) + a_h(U_h, \phi) = m_h(U_h, \partial_h^\bullet \phi)$$

follow from Banach-Nečas-Babuška theorem, in the same way as this was proved for the continuous equation [16].

4. SEMI-DISCRETIZATION IN SPACE

The next step is to consider the finite element subspace $S_h(t) \subset H^1(\Gamma_h(t))$ and to prove existence and uniqueness of the solution of a semi-discrete problem. For the proof of well-posedness we consider the following Gelfand triple

$$(23) \quad S_h(t) \subset L_h(t) \subset S_h^*(t),$$

where the spaces are the same as sets and they are defined by (19), but they have different norms which they inherit from corresponding continuous spaces, i.e.

$$S_h(t) := (S_h(t), \|\cdot\|_{H^1(\Gamma_h(t))}) \quad \text{and} \quad L_h(t) := (S_h(t), \|\cdot\|_{L^2(\Gamma_h(t))}).$$

The dual space $S_h^*(t)$ is defined as a set of all continuous linear functionals on $S_h(t)$ with the standard dual norm (see for example [40]) defined by $\|v\|_{S_h^*(t)} := \sup_{u \in B(S_h)} \|v(u)\|$. Note that all three norms are equivalent as norms on finite dimensional spaces, which implies that (23) is the Gelfand triple. Hence, the Gelfand triple for our semi-discrete problem will be

$$(24) \quad L^2(\Omega, S_h(t)) \subset L^2(\Omega, L_h(t)) \subset L^2(\Omega, S_h^*(t)).$$

Setting

$$V_h(t) := L^2(\Omega, S_h(t)) \quad \text{and} \quad H_h(t) := L^2(\Omega, L_h(t)),$$

we can define the solution space for the semi-discrete problem as a space of the functions $\phi_h(t) \in V_h(t)$ that are smooth for each path, for all $\omega \in \Omega$, $\phi_h(\omega) \in \mathcal{C}_{S_h}^1$ i.e. $\partial_h^\bullet \phi_h$ is defined path-wise for path-wise smooth functions and in addition $\partial_h^\bullet \phi_h(t) \in H_h(t)$

$$W_h(V_h, H_h) := \{\phi_h \in L_{V_h}^2 \mid \omega \mapsto \phi_h(\omega) \in \mathcal{C}_{S_h}^1 \text{ and } \partial_h^\bullet \phi_h(t) \in H_h(t)\}.$$

The norm of the space is defined by

$$\|\phi_h^2\|_{W_h(V_h, H_h)} := \int_0^T \|\phi_h(t)\|_{L^2(\Omega, H^1(\Gamma_h(t)))}^2 + \|\partial_h^\bullet \phi_h(t)\|_{L^2(\Omega, L^2(\Gamma_h(t)))}^2$$

Definition 2. *The semi-discrete in space problem is:*

Find $U_h \in W_h(V_h, H_h)$ such that $U_h(0) = U_{h,0} \in V_h(0)$ and

$$(25) \quad \frac{d}{dt} m_h(U_h, \phi) + a_h(U_h, \phi) = m_h(U_h, \partial_h^\bullet \phi) \quad \text{for all } \phi \in W_h(V_h, H_h).$$

Theorem 1. *The semi-discrete problem (25) has a unique solution $U_h \in W_h(V_h, H_h)$ which satisfies*

$$\|U_h\|_{W(V_h, H_h)} \leq C \|U_{h,0}\|_{V_h(0)}.$$

Proof. Let $\omega \in \Omega$ be arbitrary. From the deterministic results [20, Lemma 4.3] we know that there exists a unique solution $U_h(\omega) \in D_h := \{\phi_h, \partial_h^\bullet \phi_h \in \mathcal{C}^0(\mathcal{G}_T^h) \mid \phi_h(t) \in S_h(t), \text{ for all } t\}$ such that for all $\varphi_h \in D_h$ and $t \in (0, T]$

$$(26) \quad \frac{d}{dt} \int_{\Gamma_h(t)} U_h(\omega) \varphi_h + \int_{\Gamma_h(t)} \alpha^{-l} \nabla_{\Gamma_h} U_h(\omega) \nabla_{\Gamma_h} \varphi_h = \int_{\Gamma_h} U_h(\omega) \partial_h^\bullet \varphi_h$$

and $U_h(\omega; 0) = U_{h,0}(\omega)$. Hence, $U_h(\omega)$ is a path-wise solution. The next step is to prove that $U_h(t) \in V_h(t)$. We start with proving the measurability of the map $\omega \mapsto U_h(\omega, t)$. Since for every t we have that $U_h(\omega, t) \in S_h(t)$, we can write

$$(27) \quad U_h(\omega, t, x) = \sum_{j=1}^N U_j(\omega, t) \phi_j(x, t)$$

Inserting (27) into the equation (26) and taking $\varphi_h = \phi_i, i = 1, \dots, N$ we get

$$(28) \quad \frac{d}{dt} \sum_{j=1}^N U_j(\omega) \int_{\Gamma_h(t)} \phi_j \phi_i + \sum_{j=1}^N U_j(\omega) \int_{\Gamma_h(t)} \alpha^{-l}(\omega) \nabla_{\Gamma_h} \phi_j \nabla_{\Gamma_h} \phi_i = 0$$

where we also used the transport property (21) of the basis functions. Setting $U(\omega, t) = [U_j(\omega, t)]$, the evolving mass matrix

$$M(t)_{ij} := \int_{\Gamma_h(t)} \phi_i(t) \phi_j(t)$$

and the evolving stiffness matrix

$$S(\omega, t)_{ij} := \int_{\Gamma_h(t)} \alpha^{-l}(\omega, t) \nabla_{\Gamma_h} \phi_j(t) \nabla_{\Gamma_h} \phi_i(t)$$

we can write (28) in the matrix form as

$$M(t) \frac{\partial}{\partial t} U(\omega, t) + (M'(t) + S(\omega, t)) U(\omega, t) = 0, \quad U(\omega, 0) = U_0(\omega).$$

Solving the previous matrix equation we obtain

$$U(\omega, t) = \exp \left(- \int_0^T \chi_{[0,T]}(s) M^{-1}(s) (M'(s) + S(\omega, s)) dt \right) U_0(\omega).$$

Since α^{-l} is an $\mathcal{F} \otimes \mathcal{B}(\mathcal{G}_T^h)$ -measurable function by Fubini's Theorem [25, Sec. 36, Thm. C] it follows that

$$\omega \mapsto S(\omega, s)$$

is $\mathcal{F}$ -measurable. In a similar way we obtain that the map

$$\omega \mapsto \int_0^T \chi_{[0,T]}(s) M^{-1}(s) \left(\frac{d}{dt} M(s) + S(\omega, s) \right) dt$$

is $\mathcal{F}$ -measurable. From this conclusion, together with the continuity of the exponential map and $\mathcal{F}$ -measurability of the initial function $U_{h,0}$, it follows that $\omega \mapsto U_h(\omega)$ is $\mathcal{F}$ -measurable.

Furthermore, from our assumptions on the time regularity of the domain, we see that the matrices M , $S(\omega, \cdot)$ and M^{-1} are $\mathcal{C}^1(0, 1)$ functions thus we can directly read that the coefficients $U(\omega, \cdot)$ are $\mathcal{C}^1(0, T)$. This implies that $U_h(\omega, \cdot, \cdot) \in \mathcal{C}_{S_h}^1$.

The next step is to prove the boundedness of the $L^2(\Omega)$ norm of the solution U_h . This bound follows directly from the path-wise stability results from [20, Lemma 4.3] which implies

$$\|U_h(t, \omega)\|_{L^2(\Gamma_h(t))}^2 + \|\nabla_{\Gamma_h} U_h(\omega, t)\|_{L^2(\Gamma_h(t))}^2 \leq C \|U_{h,0}(\omega)\|_{L^2(\Gamma_h)}$$

Since $U_{h,0}(x) \in L^2(\Omega)$ and C depends just on the geometry of the $\Gamma_h(t)$ and $U_{h,0}$ it is independent on ω , it follows that $U_h(t) \in V_h(t)$.

Hence, $U_h \in W_h(V_h, H_h)$. It is left to show that U_h is a solution of semi-discrete scheme (25) for appropriate test space, i.e. we know that $U_h(\omega)$ solves (26) and with measurability results that we proved, we can integrate this relation over Ω , however it is left to show that the relation will still hold for every test function from $W_h(V_h, H_h)$. The relation (25) holds for a function $\phi_h(\omega, x, t) := \eta(\omega) \varphi_h(x, t)$, $\eta \in L^2(\Omega)$, $\varphi_h \in W_h(S_h, L_h)$. Since $W_h(V_h, H_h) \cong L^2(\Omega) \otimes W_h(S_h, L_h)$, and

$$\{\varphi_h(x, t) \eta(\omega) \mid \varphi_h \in W_h(S_h, L_h), \eta \in L^2(\Omega)\} \subset L^2(\Omega) \otimes W_h(S_h, L_h)$$

is dense, it follows that (25) holds for every $\phi \in W_h(V_h, H_h)$. This completes the proof. $\square$

Let U_h be the solution of the semi-discrete scheme (25) and let $u_h = U_h^l$ be its lift. Then, as a consequence of Lemma 1 and the expression that connects the material derivatives of U_h and u_h (see [18, Theorem 6.2]), the following estimate holds

$$(29) \quad \|u_h\|_{W(V, H)} \leq C_0 \|u_h^0\|_{L^2(\Omega, L^2(\Gamma_0))}.$$

5. APPROXIMATION ESTIMATES

In this section we will formulate the results concerning the approximation of the surface, which are in the deterministic setting proved in [18] and [20]. Our goal is to prove that they still hold in the random case. The main task is to keep track of constants that appear and show that they are independent of realization. This conclusion mainly follows from the assumption (11) about the uniform distribution of the diffusion coefficient. Furthermore, we need to show that the extended definitions of the interpolation operator and Ritz projection operator are integrable with respect to ω .

We start with an interpolation estimate for functions $\eta \in L^2(\Omega, H^2(\Gamma(t)))$, where the interpolation $I_h \eta$ is defined as a lift of a standard linear interpolation $\tilde{I}_h \eta \in L^2(\Omega, S_h(t))$. Note that the assumption that vertices of $\Gamma_h(t)$ lie on the smooth surface $\Gamma(t)$ and $n \leq 3$, provide that $\tilde{I}_h$ is well-defined.

Lemma 4. *Given $\eta \in L^2(\Omega, H^2(\Gamma(t)))$ there exists a unique $I_h \eta \in L^2(\Omega, S_h^l(t))$ such that*

$$(30) \quad \begin{aligned} & \|\eta - I_h \eta\|_{L^2(\Omega, L^2(\Gamma(t)))} + h \|\nabla_\Gamma(\eta - I_h \eta)\|_{L^2(\Omega, L^2(\Gamma(t)))} \\ & \leq ch^2(\|\nabla_\Gamma^2 \eta\|_{L^2(\Omega, L^2(\Gamma(t)))} + h \|\nabla_\Gamma \eta\|_{L^2(\Omega, L^2(\Gamma(t)))}). \end{aligned}$$

Proof. The proof of the lemma follows directly from the deterministic case and Lemma 1. $\square$

We continue with bounding the geometric perturbation errors in the bilinear forms.

Lemma 5. *$W_h(\cdot, t), \phi_h(\cdot, t) \in L^2(\Omega, S_h(t))$ with corresponding lifts $w_h(\cdot, t), \varphi_h(\cdot, t) \in L^2(\Omega, S_h^l(t))$ the following bounds hold*

$$(31) \quad |m(w_h, \varphi_h) - m_h(W_h, \phi_h)| \leq ch^2 \|w_h\|_{L^2(\Omega, L^2(\Gamma))} \|\varphi_h\|_{L^2(\Omega, L^2(\Gamma))}$$

$$(32) \quad |a(w_h, \varphi_h) - a_h(W_h, \phi_h)| \leq ch^2 \|\nabla_\Gamma w_h\|_{L^2(\Omega, L^2(\Gamma))} \|\nabla_\Gamma \varphi_h\|_{L^2(\Omega, L^2(\Gamma))}$$

$$(33) \quad |g(w_h; w_h, \varphi_h) - g_h(W_h; W_h, \phi_h)| \leq ch^2 \|w_h\|_{L^2(\Omega, H^1(\Gamma))} \|\varphi_h\|_{L^2(\Omega, H^1(\Gamma))}.$$

$$(34) \quad |m(\partial_h^\bullet w_h, \varphi_h) - m_h(\partial_h^\bullet W_h, \phi_h)| \leq ch^2 \|\partial_h^\bullet w_h\|_{L^2(\Omega, L^2(\Gamma))} \|\varphi_h\|_{L^2(\Omega, L^2(\Gamma))}.$$

Proof. The results follow from the the uniform bound of α and $\partial_h^\bullet \alpha$, and analog deterministic results, that can be found [20] and [33]. $\square$

Since the velocity is deterministic, we can use the result from [20, Lemma 5.6] about the difference between the continuous and discrete velocity on the smooth surface Γ . Furthermore, its corollary gives us the estimate for the difference between continuous and discrete material derivative (see [20, Corollary 5.7]).

Lemma 6. *Let v be the continuous velocity and v_h the discrete velocity on $\Gamma(t)$. Then we have the following estimates*

$$(35) \quad |v - v_h| + h \|\nabla_\Gamma(v - v_h)\| \leq ch^2$$

$$(36) \quad \|\partial^\bullet z - \partial_h^\bullet z\|_{H(t)} \leq ch^2 \|z\|_{V(t)} \quad z \in V(t)$$

$$(37) \quad \|\nabla_\Gamma(\partial^\bullet z - \partial_h^\bullet z)\|_{H(t)} \leq ch \|z\|_{L^2(\Omega, H^2(\Gamma))} \quad z \in L^2(\Omega, H^2(\Gamma(t))),$$

when the left hand sides are well-defined.

5.1. Ritz projection and error analysis. In this section we will define the Ritz projection u_p of the solution u , which we will use later in order to estimate the error between u and semi-discrete solution $u_h := U_h^l$. Furthermore we will give results about the error estimates for the projection.

We start with the standard path-wise definition of the Ritz projection for each ω and t . Since $S_h^l(t)$ is a finite dimensional subset of $H^1(\Gamma(t))$, for every $v \in H^1(\Gamma(t))$ such that $\int_{\Gamma(t)} v(x) = 0$, there exists unique $\mathcal{R}_{h,t}^{\alpha(\omega)} v \in S_h^l$ such that $\int_{\Gamma(t)} \mathcal{R}_{h,t}^{\alpha(\omega)} v = 0$ and

$$(38) \quad \int_{\Gamma(t)} \alpha(\omega) \nabla_\Gamma v \cdot \nabla_\Gamma \varphi_h = \int_{\Gamma(t)} \alpha(\omega) \nabla_\Gamma \mathcal{R}_{h,t}^{\alpha(\omega)} v \cdot \nabla_\Gamma \varphi_h \quad \forall \varphi_h \in S_h^l(t).$$

Note that in order to simplify the notation we omit writing the time dependence of the coefficient α , but instead just write $\alpha(\omega)$. We will also make use of the function $\mathcal{R}_{h,t}^\beta$ and note that this is well defined for $\beta \in L^\infty(\Gamma(t))$ such that there exists constants $\beta_{\min}, \beta_{\max} > 0$ such that $\beta_{\min} \leq \beta \leq \beta_{\max}$.

Since our solution is a random variable, we want to estimate the $L^2(\Omega, S_h^l(t))$ error. Therefore, we define a map u_p that maps each ω , i.e. each realization of the diffusion coefficient $\alpha(\omega)$ to the corresponding path-wise Ritz projection $\mathcal{R}_{h,t}^{\alpha(\omega)}$. In the following lemma we prove the 2-integrability of u_p with respect $\mathbb{P}$ and its a -orthogonality.

Lemma 7. *For a given $u \in L^2(\Omega, H^1(\Gamma(t)))$ such that $\int_{\Gamma(t)} u(\omega, x) = 0$ for every ω , the function u_p defined by*

$$(39) \quad u_p(\omega) := \mathcal{R}_{h,t}^{\alpha(\omega)}(u(\omega))$$

belongs to $L^2(\Omega, S_h^l)$. Furthermore, u_p is unique and it satisfies

$$(40) \quad a(u, \eta_h) = a(u_p, \eta_h), \quad \forall \eta_h \in V_h(t).$$

Proof. We begin by proving the measurability of the map $\Omega \ni \omega \mapsto u_p(\omega) \in H^1(\Gamma(t))$. We consider a map $R_{h,t} : L^\infty(\mathcal{G}_T) \rightarrow \mathcal{L}(H^1(\Gamma(t)), S_h^l(t))$ that is a path-wise Ritz projection as a function of the diffusion coefficient defined by $R_{h,t} := \mathcal{R}_{h,t}^{\alpha(\omega)}$ and prove that it is continuous. Let $(\alpha_n)_n$ and α be from $L^\infty(\mathcal{G}_T)$ such that $\alpha_n \rightarrow \alpha$, as $n \rightarrow \infty$. We may assume that $0 < \alpha_{\min}/2 \leq \alpha_n \leq 2\alpha_{\max}$ and $\mathcal{R}_{h,t}^{\alpha_n}$ is a well defined operator for each n . Using the uniform bound (11) and relation (38) with the test function $\varphi_h = (\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha)u \in S_h^l(t)$ we obtain

$$\begin{aligned} \alpha_{\min}^2 \quad & \|(\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha)u\|_{H^1(\Gamma)}^2 \leq C_P^2 \int_{\Gamma} \alpha |\nabla_{\Gamma}(\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha)u|^2 \\ &= C_P^2 \left(\int_{\Gamma} (\alpha - \alpha_n) \nabla_{\Gamma} \mathcal{R}_{h,t}^{\alpha_n} u \nabla_{\Gamma} (\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha) u \right. \\ &+ \left. \int_{\Gamma} \alpha_n \nabla_{\Gamma} \mathcal{R}_{h,t}^{\alpha_n} u \nabla_{\Gamma} (\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha) u - \int_{\Gamma} \alpha \nabla_{\Gamma} u \nabla_{\Gamma} (\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha) u \right) \\ &= C_P^2 \left(\int_{\Gamma} (\alpha_n - \alpha) (\nabla_{\Gamma} u - \nabla_{\Gamma} \mathcal{R}_{h,t}^{\alpha_n} u) \nabla_{\Gamma} (\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha) u \right) \\ &\leq C_P^2 \|\alpha_n - \alpha\|_{\infty} \|u - \mathcal{R}_{h,t}^{\alpha_n} u\|_{H^1(\Gamma)} \|(\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha)u\|_{H^1(\Gamma)} \\ &\leq 2C_P^2 \|\alpha_n - \alpha\|_{\infty} \|u\|_{H^1(\Gamma)} \|(\mathcal{R}_{h,t}^{\alpha_n} - \mathcal{R}_{h,t}^\alpha)u\|_{H^1(\Gamma)} \end{aligned}$$

where C_P is the Poincaré constant. The last inequality implies the continuity of the map $R_{h,t}$. Therefore, the map u_p is a composition of a measurable map $\alpha : \Omega \rightarrow L^\infty(\mathcal{G}_T)$ and a continuous map $R_{h,t}$. Thus by, e.g., [26, Lemma A.5] u_p is measurable.

We next prove the boundedness of the norm. Let $u \in L^2(\Omega, H^1(\Gamma(t)))$ such that $\int_{\Gamma(t)} u(\omega, x) = 0$ almost everywhere. Then for every ω there exists a unique $u_p(\omega) := \mathcal{R}_{h,t}^{\alpha(\omega)}(u(\omega)) \in S_h^l(t)$ such that (38) holds. Combining this with Poincaré's inequality [22, Theorem 2.12] and (11) we obtain

$$\begin{aligned} \alpha_{\min} \int_{\Omega} \|u_p(\omega)\|_{H^1(\Gamma(t))}^2 &\leq C \int_{\Omega} \alpha(\omega) \|\nabla_{\Gamma} \mathcal{R}_{h,t}^{\alpha(\omega)}(u(\omega))\|_{L^2(\Gamma(t))}^2 \\ &\leq C \alpha_{\max} \int_{\Omega} \|\nabla_{\Gamma} u(\omega)\|_{L^2(\Gamma(t))}^2 \leq C \|\nabla_{\Gamma} u\|_{L^2(\Omega, L^2(\Gamma(t)))}^2, \end{aligned}$$

which implies $u_p \in L^2(\Omega, S_h^l(t))$.

It is left to show that if we take integral over Ω of (38), equality will still hold for every test function $\eta_h \in V_h(t)$. Let us first take the test function of the form $\eta_h(\omega, x) = \varphi_h(x)w(\omega)$, $\varphi_h \in S_h^l(t)$, $w \in L^2(\Omega)$. From (38) we get

$$\begin{aligned} & \int_{\Omega} \int_{\Gamma(t)} \alpha(\omega, x) \nabla_{\Gamma} u(\omega, x) \nabla_{\Gamma} (\varphi_h(x)w(\omega)) = \\ & \int_{\Omega} w(\omega) \int_{\Gamma(t)} \alpha(\omega, x) \nabla_{\Gamma} u_p(\omega, x) \nabla_{\Gamma} \varphi_h(x) = \\ & \int_{\Omega} \int_{\Gamma(t)} \alpha(\omega, x) \nabla_{\Gamma} u_p(\omega, x) \nabla_{\Gamma} (\varphi_h(x)w(\omega)). \end{aligned}$$

Since $V_h(t) \cong L^2(\Omega) \otimes S_h^l(t)$ and $\{v(x)w(\omega) | v \in S_h^l(t), w \in L^2(\Omega)\}$ is a dense subset of $L^2(\Omega) \otimes S_h^l(t)$, the relation (40) follows. $\square$

The error in the projection is established by the next theorem.

Theorem 2. For $u \in L^2(\Omega, H^2(\Gamma(t)))$ the error in the projection satisfies

$$(41) \quad \|u - u_p\|_{L^2(\Omega, L^2(\Gamma(t)))} + h \|\nabla_{\Gamma}(u - u_p)\|_{L^2(\Omega, L^2(\Gamma(t)))} \leq ch^2 \|u\|_{L^2(\Omega, H^2(\Gamma(t)))}.$$

Proof. The L^2 -bound for the gradient follows from the standard error estimates for the surface finite element method (see [17]) that uses the error for the gradient of the interpolation (30) and in addition we use the uniform bound for α .

In order to get L^2 -estimate we will use Aubin-Nitsche duality argument. The path-wise dual problem is: for every $\omega \in \Omega$ find $w \in H^2(\Gamma(t))$ with $\int_{\Gamma(t)} w = 0$ such that

$$-\nabla_{\Gamma} \cdot (\alpha \nabla_{\Gamma} w) = u - u_p.$$

Then from the standard results about the random elliptic PDEs with uniformly bounded coefficients (see [11, 8]) it is known that $w \in L^2(\Omega, H^1(\Gamma(t)))$, thus the map $\omega \rightarrow w(\omega)$ is measurable. Furthermore, since in addition Γ is $\mathcal{C}^2$, by [11, Prop. 3.1] it follows that $w(\omega) \in H^2(\Gamma(t))$ with the bound

$$(42) \quad \|w(\omega)\|_{H^2(\Gamma(t))} \leq \frac{\alpha_{\max}}{\alpha_{\min}^3} \|\alpha(\omega)\|_{\mathcal{C}^1(\bar{\Gamma}(t))} \|u - u_p\|_{L^2(\Gamma(t))}.$$

By Assumption 1 and Assumption 2, the right hand side of (42) is uniformly bounded and we obtain

$$\|w\|_{L^2(\Omega, H^2(\Gamma(t)))} \leq c \|u - u_p\|_{L^2(\Omega, L^2(\Gamma(t)))}.$$

Using previous estimate, (30) and (40), we get

$$\begin{aligned} \|u - u_p\|_{L^2(\Omega, L^2(\Gamma(t)))}^2 &= a(w, u - u_p) = a(w - I_h w, u - u_p) \\ &\leq \alpha_{\max} \|\nabla_{\Gamma}(w - I_h w)\|_{L^2(\Omega, L^2(\Gamma(t)))} \|\nabla_{\Gamma}(u - u_p)\|_{L^2(\Omega, L^2(\Gamma(t)))} \\ &\leq ch \|w\|_{L^2(\Omega, H^2(\Gamma(t)))} \|\nabla_{\Gamma}(u - u_p)\|_{L^2(\Omega, L^2(\Gamma(t)))} \\ &\leq ch^2 \|u - u_p\|_{L^2(\Omega, L^2(\Gamma(t)))} \|u\|_{L^2(\Omega, H^2(\Gamma(t)))}. \end{aligned}$$

Thus,

$$(43) \quad \|u - u_p\|_{L^2(\Omega, L^2(\Gamma(t)))} \leq ch^2 \|u\|_{L^2(\Omega, H^2(\Gamma(t)))}$$

which completes the proof of the theorem. $\square$

Note that without the Assumption 2 we would still get the $H^1(\Gamma(t))$ bound. We state the analogue to the result from the deterministic setting [20, Theorem 6.2] about the estimate for the error in the material derivative of u_p .

Theorem 3. *The error in the material derivative of the Ritz projection satisfies the bounds*

$$(44) \quad \begin{aligned} & \|\partial_h^\bullet u - \partial_h^\bullet u_p\|_{L^2(\Omega, L^2(\Gamma))} + h \|\nabla_\Gamma(\partial_h^\bullet u - \partial_h^\bullet u_p)\|_{L^2(\Omega, L^2(\Gamma))} \\ & \leq ch^2(\|u\|_{L^2(\Omega, H^2(\Gamma))} + \|\partial^\bullet u\|_{L^2(\Omega, H^2(\Gamma))}). \end{aligned}$$

5.2. Error bound for the ESFEM.

Theorem 4. *Let u be sufficiently smooth solution of (10) that satisfies*

$$(45) \quad \int_0^T \|u\|_{L^2(\Omega, H^2(\Gamma(t)))}^2 + \|\partial^\bullet u\|_{L^2(\Omega, H^2(\Gamma(t)))}^2 dt < \infty$$

and let $u_h(\cdot, \cdot, t) := U_h^l(\cdot, \cdot, t)$, $t \in [0, T]$ with initial data $u_{h0} = U_{h0}^l$ that satisfies

$$(46) \quad \|u(\cdot, \cdot, 0) - u_{h0}\|_{L^2(\Omega, L^2(\Gamma_0))} \leq ch^2.$$

Then the error estimate

$$(47) \quad \sup_{t \in (0, T)} \|u(\cdot, \cdot, t) - u_h(\cdot, \cdot, t)\|_{L^2(\Omega, L^2(\Gamma(t)))} \leq ch^2$$

holds for deterministic constant c that is independent on h but depends on norms (45).

Proof. Let us note that using the error analysis from the previous sections we can now proceed analogously to the proof of L^2 -estimate for ESFEM [20, Theorem 4.4]. The important remark for our probabilistic setting is that these estimates will stay deterministic, since all previous error bounds are deterministic. For the completeness of the paper, we will present the main steps in the proof.

The basic step is to decompose the error into errors between the smooth solution and Ritz projection and Ritz projection and discrete solution:

$$u - u_h = u - u_p + u_p - u_h.$$

Theorem 2 states the estimate for $u - u_p$, thus it remains to estimate the norm of

$$\theta := u_p - u_h \in L^2(\Omega, S_h^l).$$

First step is to derive the equation for θ . Subtracting equation for u_p and u_h we get

$$(48) \quad \frac{d}{dt} m(\theta, \varphi_h) + a(\theta, \varphi_h) - m(\theta, \partial_h^\bullet \varphi_h) = F_2(\varphi_h) - F_1(\varphi_h), \quad \forall \varphi_h \in L^2(\Omega, S_h^l).$$

where

$$(49) \quad \begin{aligned} F_1(\varphi_h) &= m(\partial_h^\bullet u_h, \varphi_h) - m(\partial_h^\bullet U_h, \phi_h) \\ &+ a(u_h, \varphi_h) - a_h(U_h, \phi_h) + g(v_h; u_h, \varphi_h) - g(V_h; U_h, \phi_h) \end{aligned}$$

$$(50) \quad F_2(\varphi_h) = m(u, \partial^\bullet \varphi_h - \partial_h^\bullet \varphi_h) + m(\rho, \partial_h^\bullet \varphi_h) - \frac{d}{dt} m(\rho, \varphi_h).$$

Integrating (48) in time, testing it with $\varphi = \theta$ and using partial integration and transport theorem we get

$$\frac{1}{2} \|\theta(t)\|_{L^2(\Omega, L^2(\Gamma))}^2 - \frac{1}{2} \|\theta(0)\|_{L^2(\Omega, L^2(\Gamma_0))}^2 + \int_0^t a(\theta, \theta) + \int_0^t g(v_h; \theta, \theta) = \int_0^t F_2(\theta) - F_1(\theta).$$

Estimates of $F_1(\theta)$ and $F_2(\theta)$ follow from Lemma 5, errors of projection (41) and (44) and Lemma 6

$$\begin{aligned} |F_1(\theta)| &\leq ch^2 \|\partial_h^\bullet u_h\|_{L^2(\Omega, L^2(\Gamma))} \|\theta_h\|_{L^2(\Omega, L^2(\Gamma))} + ch^2 \|u_h\|_{L^2(\Omega, H^1(\Gamma))} \|\theta_h\|_{L^2(\Omega, H^1(\Gamma))} \\ |F_2(\theta)| &\leq ch^2 (\|u\|_{L^2(\Omega, H^2(\Gamma))} + \|\partial^\bullet u\|_{L^2(\Omega, H^2(\Gamma))}) \|\theta_h\|_{L^2(\Omega, L^2(\Gamma))} + \\ &\quad ch^2 \|u\|_{L^2(\Omega, L^2(\Gamma))} \|\theta_h\|_{L^2(\Omega, H^1(\Gamma))}. \end{aligned}$$

Applying these estimates, uniform condition and Young's inequality to (5.2) we arrive at

$$\begin{aligned} &\frac{1}{2} \|\theta(t)\|_{L^2(\Omega, L^2(\Gamma))}^2 + (\alpha_{max} - \varepsilon) \int_0^t \|\nabla_\Gamma \theta\|_{L^2(\Omega, L^2(\Gamma))}^2 \\ &\leq \frac{1}{2} \|\theta(0)\|_{L^2(\Omega, L^2(\Gamma_0))}^2 + c(\varepsilon) \int_0^t \|\theta\|_{L^2(\Omega, L^2(\Gamma))}^2 \\ &+ c(\varepsilon) h^4 \int_0^t [\|u\|_{L^2(\Omega, H^2(\Gamma))}^2 + \|\partial^\bullet u\|_{L^2(\Omega, H^2(\Gamma))}^2 + \|\partial_h^\bullet u\|_{L^2(\Omega, L^2(\Gamma))}^2 + \|u_h\|_{L^2(\Omega, H^1(\Gamma))}^2]. \end{aligned}$$

For small enough ε , Gronwall's inequality implies

$$(51) \quad \sup_{(0, T)} \|\theta(t)\|_{L^2(\Omega, L^2(\Gamma))}^2 + \int_0^T \|\nabla_\Gamma \theta\|_{L^2(\Omega, L^2(\Gamma))}^2 \leq c \|\theta(0)\|_{L^2(\Omega, L^2(\Gamma(0)))}^2 + ch^4 C,$$

where

$$C = \int_0^T [\|u\|_{L^2(\Omega, H^2(\Gamma))}^2 + \|\partial^\bullet u\|_{L^2(\Omega, H^2(\Gamma))}^2 + \|\partial_h^\bullet u\|_{L^2(\Omega, L^2(\Gamma))}^2 + \|u_h\|_{L^2(\Omega, H^1(\Gamma))}^2].$$

It is left to derive the bound on $\theta(0)$ and to prove that C is deterministic and does not depend on h . The bound on $\theta(0)$ follows from the assumption (46). To prove the deterministic bound on C we use assumption (45) and a priori bound on the lifted solution (29), which completes the proof. $\square$

Remark 5. Observe that without Assumption 2 we still get the H^1 -bound

$$\int_0^T \|\nabla_\Gamma(u(t) - u_h(t))\|_{L^2(\Omega, L^2(\Gamma(t)))}^2 \leq ch^2.$$

6. SINGLE-LEVEL MONTE-CARLO METHOD FOR PARABOLIC RANDOM PDES ON MOVING HYPERSURFACE

As already discussed, since the solution of considered random PDE is a random variable, we are interested in approximation of its expectation. In this section we deal with uncertainty using (single-level) Monte Carlo (MC). Using the error estimate from Theorem 4, we derive a result on error bound of MC-ESFEM.

We start with recalling some basic results about Monte Carlo. Let $Y \in L^2(\Omega, H)$, where H is separable Hilbert space. Then the Monte Carlo estimator of Y is defined by

$$E_M[Y] := \frac{1}{M} \sum_{i=1}^M \hat{Y}^i \quad M \in \mathbb{N},$$

where $\hat{Y}^i$ are independent identically distributed copies of a random variable Y . Note that $E_M[Y]$ is also a random variable on $\Omega^M := \bigotimes_{i=1}^M \Omega$.

Lemma 8. *For any $M \in \mathbb{N}$ and $Y \in L^2(\Omega, H)$, it holds that*

$$(52) \quad \|E[Y] - E_M[Y]\|_{L^2(\Omega^M, H)} = \frac{1}{\sqrt{M}} \text{Var}[Y]^{\frac{1}{2}} \leq \frac{1}{\sqrt{M}} \|Y\|_{L^2(\Omega, H)},$$

where $\text{Var}[Y] = E[\|E[Y] - Y\|_H^2]$.

Proof. This is the general result for the Single-level Monte Carlo approximation and the proof can be found for example in [4, Lemma 4.1]. $\square$

We will use previous lemma to estimate the Monte Carlo error of the semi-discrete solution u_h .

Theorem 5. *Under the same assumptions as in Theorem 3, the following error bound holds*

$$\sup_{t \in (0, T)} \|E[u(t)] - E_M[u_h(t)]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} \lesssim h^2 + \frac{1}{\sqrt{M}} h^2 + \frac{1}{\sqrt{M}}.$$

Proof. Directly from the Cauchy-Schwarz inequality it follows

$$(53) \quad \|E[u]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} \leq \|u\|_{L^2(\Omega^M, L^2(\Gamma(t)))}$$

Using the previous inequality and (52), we arrive at

$$(54) \quad \begin{aligned} & \|E[u(t)] - E_M[u_h(t)]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} \\ & \leq \|E[u(t)] - E[u_h(t)]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} + \|E[u_h(t)] - E_M[u_h(t)]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} \\ & \leq \|u(t) - u_h(t)\|_{L^2(\Omega, L^2(\Gamma))} + M^{-1/2} \|u_h(t)\|_{L^2(\Omega, L^2(\Gamma))}. \end{aligned}$$

Furthermore, L^2 -error estimate from Theorem 3 gives us

$$\begin{aligned} \|u_h(t)\|_{L^2(\Omega, L^2(\Gamma(t)))} & \leq \|u(t) - u_h(t)\|_{L^2(\Omega, L^2(\Gamma(t)))} + \|u(t)\|_{L^2(\Omega, L^2(\Gamma(t)))} \\ & \leq ch^2 (\|u(t)\|_{L^2(\Omega, H^2(\Gamma(t)))} + 1) + \|u(t)\|_{L^2(\Omega, L^2(\Gamma(t)))}. \end{aligned}$$

Combining previous estimate with (54) we get

$$\begin{aligned} & \sup_{t \in (0, T)} \|E[u(t)] - E_M[u_h(t)]\|_{L^2(\Omega^M, L^2(\Gamma(t)))} \leq ch^2 \left(\sup_{t \in (0, T)} \|u(t)\|_{L^2(\Omega, H^2(\Gamma(t)))} + 1 \right) + \\ & \frac{1}{\sqrt{M}} ch^2 \left(\sup_{t \in (0, T)} \|u(t)\|_{L^2(\Omega, H^2(\Gamma(t)))} + 1 \right) + \frac{1}{\sqrt{M}} \sup_{t \in (0, T)} \|u(t)\|_{L^2(\Omega, H^2(\Gamma(t)))} = \\ & ch^2 + c \frac{1}{\sqrt{M}} h^2 + c \frac{1}{\sqrt{M}}, \end{aligned}$$

which proves the theorem. $\square$

7. NUMERICAL EXPERIMENTS

The above methods were discretised in time using a backward Euler method; see [21] for details and further numerical analysis in the deterministic case. The result is at each time step and each sample a system of linear equations for the nodal values of a finite element function which is solved using a conjugate gradient linear solver.

The above method is methods were implemented using the Distributed Unified Numerics Environment (DUNE) [5, 6, 14].

The code produced to run these computations is available at

<https://github.com/tranner/dune-mcesfem>

7.1. Example on a moving curve. We take a curve defined by the zero level set of the function

$$\Phi(x, t) = \frac{x_1^2}{a(t)} + \frac{x_2^2}{b(t)},$$

where $a(t) = 1 + \frac{1}{4} \sin(t)$ and $b(t) = 1 + \frac{1}{4} \cos(t)$. We take (9) and parametrise the diffusion coefficient and right hand side using uniformly distributed random variables $Y_1, Y_2 \sim \mathcal{U}(-1, 1)$. We consider a parametric stochastic diffusion coefficient α to be given by

$$\alpha(x, \omega) = 1 + \frac{Y_1}{4} \sin(2x_1) + \frac{Y_2}{4} \sin(2x_2)$$

and right hand side function f so that the exact solution of the resulting pathwise problem is given by

$$u(x, t, \omega) = \sin(t) \{ \cos(3x_1) + \cos(3x_2) + Y_1 \cos(5x_1) + Y_2 \cos(5x_2) \}.$$

Finally we set $u_0 = 0$.

We discretise as above and take 20 i.i.d. samples of the norm of the errors

$$\|\mathbb{E}u - \mathbb{E}_M u_h^\ell\|_{L^\infty(0, T; L^2(\Gamma))}$$

and compute a sample average. In order to demonstrate the optimal rate of convergence between each level, two global bisectional refinements are performed, which approximately half the mesh size, the time step is reduced by a factor of four and M is increased by a factor of 16. A table of the results demonstrating the optimal scaling of the methods is shown in Table 1. The results show the $o(h^2) + o(M^{-1/2})$ rate of convergence. Errors for different ratios of M and h (with $\tau = o(h^2)$) are shown in Figure 1.

h	M	Error	eoc(h)	eoc(M)
1.50000	1	3.00350	—	—
$8.43310 \cdot 10^{-1}$	16	$2.23278 \cdot 10^{-1}$	4.51325	-0.93743
$4.34572 \cdot 10^{-1}$	256	$1.86602 \cdot 10^{-1}$	0.27066	-0.06472
$2.18962 \cdot 10^{-1}$	4096	$4.88096 \cdot 10^{-2}$	1.95642	-0.48368
$1.09692 \cdot 10^{-1}$	65536	$1.29667 \cdot 10^{-2}$	1.91768	-0.47809

TABLE 1. Results of 2d problem

7.2. Example on a moving surface. We take a surface define to be the zero level set of the function

$$\Phi(x, t) = \frac{x_1^2}{a(t)} + x_2^2 + x_3^2 - 1,$$

where $a(t) = 1 + \frac{1}{4} \sin(t)$. We take (9) with the diffusion coefficient given uniformly distributed random variables $Y_1, Y_2 \sim \mathcal{U}(-1, 1)$. We consider parametric stochastic diffusion coefficient α to be given by

$$\alpha(x, \omega) = 1 + x_1^2 + Y_1(\omega)x_1^4 + Y_2(\omega)x_2^4$$




FIGURE 1. Errors for two dimensional example. Coloured dots indicate the error given parameter values of M, h and the time step. One standard deviation about the mean is given by the grey error bars. The optimal scaling results shown in Table 1 are shown in red.

and right hand side function f so that the exact solution of the resulting pathwise problem is given by

$$u(x, t, \omega) = \sin(t)x_1x_2 + Y_1(\omega)\sin(2t)x_1^2 + Y_2(\omega)\sin(2t)x_2.$$

Finally we set $u_0 = 0$.

We discretise as above and take 8 i.i.d. samples of the norm of the errors

$$\|\mathbb{E}u - \mathbb{E}_M u_h^\ell\|_{L^\infty(0,T;L^2(\Gamma))}$$

and compute a sample average. The same scaling of h, M and time step are used as in the stationary curve example. A table of the results demonstrating the optimal scaling of the methods is shown in Table 2. Errors for different ratios of M and h (with $\tau = o(h^2)$) are shown in Figure 2.

h	M	Error	eoc(h)	eoc(M)
1.27687	1	$9.91189 \cdot 10^{-1}$	—	—
$8.31246 \cdot 10^{-1}$	16	$1.70339 \cdot 10^{-1}$	4.10285	-0.63519
$4.40169 \cdot 10^{-1}$	256	$4.61829 \cdot 10^{-2}$	2.05293	-0.47075
$2.22895 \cdot 10^{-1}$	4096	$1.18779 \cdot 10^{-2}$	1.99561	-0.48977

TABLE 2. Results of 3d problem

REFERENCES

- [1] A. Alphonse, Ch. M. Elliott, and B. Stinner. An abstract framework for parabolic PDEs on evolving spaces. *Portugaliae Mathematica*, 72(1):1–46, 2015. doi: 10.4171/pm/1955. URL <http://dx.doi.org/10.4171/PM/1955>.




FIGURE 2. Errors for three dimensional example. Coloured dots indicate the error given parameter values of M, h and the time step. One standard deviation about the mean is given by the grey error bars. The optimal scaling results shown in Table 2 are shown in red.

- [2] A. Alphonse, Ch. M. Elliott, and B. Stinner. On some linear parabolic PDEs on moving hypersurfaces. *Interfaces and Free Boundaries*, 17(2):157–187, 2015. doi: 10.4171/ifb/338. URL <http://dx.doi.org/10.4171/IFB/338>.
- [3] A. Barth, Ch. Schwab, and N. Zollinger. Multi-level monte carlo finite element method for elliptic pde’s with stochastic coefficients. *Numerische Mathematik*, 119(1):123–161, april 2011. doi: 10.1007/s00211-011-0377-0. URL <http://link.springer.com/article/10.1007/s00211-011-0377-0>.
- [4] A. Barth, A. Lang, and Ch. Schwab. Multilevel monte carlo method for parabolic stochastic partial differential equations. *BIT Numerical Mathematics*, 53(1):3–27, sep 2012. doi: 10.1007/s10543-012-0401-5. URL <http://dx.doi.org/10.1007/s10543-012-0401-5>.
- [5] P. Bastian, M. Blatt, A. Dedner, C. Engwer, R. Klöforn, M. Oldberger, and O. Sander. A generic grid interface for parallel and adaptive scientific computing. part i: Abstract framework. *Comput.*, 82:103–119, 2008.
- [6] P. Bastian, M. Blatt, A. Dedner, C. Engwer, R. Klöforn, M. Oldberger, and O. Sander. A generic grid interface for parallel and adaptive scientific computing. part ii: Impelementation and tests in DUNE. *Comput.*, 82:121–138, 2008.
- [7] M. Bertalmio, G. Sapiro, L.-T.Cheng, and S. Osher. A framework for solving surface partial differential equations for computer graphics applications. *CAM Report, UCLA, Mathematics Department*, pages 00–43, 2000.
- [8] J. Charrier. Strong and weak error estimates for elliptic partial differential equations with random coefficients. *SIAM Journal on Numerical Analysis*, 50(1):216–246, jan 2012. doi: 10.1137/100800531. URL <http://dx.doi.org/10.1137/100800531>.

- [9] J. Charrier. Numerical analysis of the advection-diffusion of a solute in porous media with uncertainty. *SIAM/ASA Journal on Uncertainty Quantification*, 3(1):650–685, jan 2015. doi: 10.1137/130937457. URL <http://dx.doi.org/10.1137/130937457>.
- [10] J. Charrier, R. Scheichl, and A. L. Teckentrup. Finite element error analysis of elliptic pdes with random coefficients and its application to multilevel monte carlo methods. *SIAM J. Numer. Anal.*, 51(1):322–352, january 2013. doi: 10.1137/110853054. URL <http://dx.doi.org/10.1137/110853054>.
- [11] J. Charrier, R. Scheichl, and A. L. Teckentrup. Finite element error analysis of elliptic PDEs with random coefficients and its application to multilevel monte carlo methods. *SIAM Journal on Numerical Analysis*, 51(1):322–352, jan 2013. doi: 10.1137/110853054. URL <http://dx.doi.org/10.1137/110853054>.
- [12] A. Cliffe, M. Giles, R. Scheichl, and A. L. Teckentrup. Multilevel monte carlo methods and applications to elliptic pdes with random coefficients. *Computing and Visualization in Science*, 14(1):3–15, august 2011. doi: 10.1007/s00791-011-0160-x. URL <http://link.springer.com/article/10.1007/s00791-011-0160-x#aboutarticle>.
- [13] K. Deckelnick, G. Dziuk, and Ch. M. Elliott. Computation of geometric partial differential equations and mean curvature flow. *Acta Numerica*, 14:139–232, may 2005. doi: 10.1017/s0962492904000224. URL <http://dx.doi.org/10.1017/S0962492904000224>.
- [14] A. Dedner., R. Klöforn, M. Nolte, and M. Ohlberger. A generic interface for parallel and adaptive discretization schemes: abstraction principles and the DUNE-FEM module. *Comput.*, 90(3-4):165–196, 2010. ISSN 0010-485X. doi: 10.1007/s00607-010-0110-3. URL <http://www.springerlink.com/index/10.1007/s00607-010-0110-3>.
- [15] A. Demlow. Higher-order finite element methods and pointwise error estimates for elliptic problems on surfaces. *SIAM Journal on Numerical Analysis*, 47(2): 805–827, jan 2009. doi: 10.1137/070708135. URL <http://dx.doi.org/10.1137/070708135>.
- [16] A. Djurdjevac. Advection-diffusion equations with random coefficients on evolving hypersurfaces. Technical report, Freie Universität Berlin, 2015. URL <http://publications.mi.fu-berlin.de/1763/>.
- [17] G. Dziuk. Finite elements for the beltrami operator on arbitrary surfaces. In *Lecture Notes in Mathematics*, pages 142–155. Springer Nature, 1988. doi: 10.1007/bfb0082865. URL <http://dx.doi.org/10.1007/BFb0082865>.
- [18] G. Dziuk and Ch. M. Elliott. Finite elements on evolving surfaces. *IMA Journal of Numerical Analysis*, 27(2):262–292, sep 2006. doi: 10.1093/imanum/drl023. URL <http://dx.doi.org/10.1093/imanum/drl023>.
- [19] G. Dziuk and Ch. M. Elliott. Surface finite elements for parabolic equations. *Journal of Computational Mathematics*, 25(4):385–407, 2007. ISSN 02549409, 19917139. URL <http://www.jstor.org/stable/43693376>.
- [20] G. Dziuk and Ch. M. Elliott. L^2 -estimates for the evolving surface finite element method. *Mathematics of Computation*, 82(281):1–24, apr 2012. doi: 10.1090/s0025-5718-2012-02601-9. URL <http://dx.doi.org/10.1090/S0025-5718-2012-02601-9>.
- [21] G. Dziuk and Ch. M. Elliott. A fully discrete evolving surface finite element method. *SIAM Journal on Numerical Analysis*, 50(5):2677–2694, 2012. doi:

- 10.1137/110828642. URL <http://dx.doi.org/10.1137/110828642>.
- [22] G. Dziuk and Ch. M. Elliott. Finite element methods for surface PDEs. *Acta Numerica*, 22:289–396, apr 2013. doi: 10.1017/s0962492913000056. URL <http://dx.doi.org/10.1017/S0962492913000056>.
- [23] Ch. M. Elliott and T. Ranner. Evolving surface finite element method for the cahn-hilliard equation. *Numerische Mathematik*, 129(3):483–534, jun 2014. doi: 10.1007/s00211-014-0644-y. URL <http://dx.doi.org/10.1007/s00211-014-0644-y>.
- [24] W. Hackbusch. *Tensor Spaces and Numerical Tensor Calculus*. Springer Berlin Heidelberg, 2012. doi: 10.1007/978-3-642-28027-6. URL <http://dx.doi.org/10.1007/978-3-642-28027-6>.
- [25] P. R. Halmos. *Measure Theory*. Graduate Texts in Mathematics. Springer, 1976. URL <http://dx.doi.org/10.1007/978-90-481-3520-2>.
- [26] L. Herrmann, A. Lang, and Ch. Schwab. Numerical analysis of lognormal diffusions on the sphere. *Preprint, arXiv:1601.02500v2*, november 2016.
- [27] V. H. Hoang and Ch. Schwab. Sparse tensor galerkin discretization of parametric and random parabolic PDEs—analytic regularity and generalized polynomial chaos approximation. *SIAM Journal on Mathematical Analysis*, 45(5):3050–3083, oct 2013. doi: 10.1137/100793682. URL <http://dx.doi.org/10.1137/100793682>.
- [28] A. J. James and J. Lowengrub. A surfactant-conserving volume-of-fluid method for interfacial flows with insoluble surfactant. *Journal of Computational Physics*, 201(2):685–722, dec 2004. doi: 10.1016/j.jcp.2004.06.013. URL <http://dx.doi.org/10.1016/j.jcp.2004.06.013>.
- [29] H. Jin, A. J. Yezzi, and S. Soatto. Region-based segmentation on evolving surfaces with application to 3d reconstruction of shape and piecewise constant radiance. *European Conference on Computer Vision*, pages 114–125, 2004.
- [30] S. C. Larsson, C. Mollet, and M. Molteni. Quasi-optimality of petrov-galerkin discretizations of parabolic problems with random coefficients. *Preprint, arXiv:1604.06611*, april 2016.
- [31] C. H. Leung and M. Berzins. A computational model for organism growth based on surface mesh generation. *Journal of Computational Physics*, 188(1):75–99, jun 2003. doi: 10.1016/s0021-9991(03)00153-0. URL [http://dx.doi.org/10.1016/S0021-9991\(03\)00153-0](http://dx.doi.org/10.1016/S0021-9991(03)00153-0).
- [32] G. Lord, C. E. Powell, and T. Shardlow. *An Introduction to Computational Stochastic PDEs*. Cambridge University Press, 2014. URL <http://dx.doi.org/10.1007/978-90-481-3520-2>.
- [33] Ch. Lubich and D. Mansour. Variational discretization of wave equations on evolving surfaces. *Mathematics of Computation*, 84(292):513–542, oct 2014. doi: 10.1090/s0025-5718-2014-02882-2. URL <http://dx.doi.org/10.1090/S0025-5718-2014-02882-2>.
- [34] O. P. Le Maître and O. M. Knio. *Spectral Methods for Uncertainty Quantification*. Springer Netherlands, 2010. doi: 10.1007/978-90-481-3520-2. URL <http://dx.doi.org/10.1007/978-90-481-3520-2>.
- [35] N. R. Morrow and G. Mason. Recovery of oil by spontaneous imbibition. *Current Opinion in Colloid and Interface Science*, 6(4):321–337, august 2001. doi: 10.1016/S1359-0294(01)00100-5. URL [http://dx.doi.org/10.1016/S1359-0294\(01\)00100-5](http://dx.doi.org/10.1016/S1359-0294(01)00100-5).

- [36] F. Nobile and R. Tempone. Analysis and implementation issues for the numerical approximation of parabolic equations with random coefficients. *Int. J. Numer. Meth. Engng.*, 80(6-7):979–1006, june 2009. doi: 10.1002/nme.2656. URL <http://dx.doi.org/10.1002/nme.2656>.
- [37] M. Reed and B. Simon. *Methods of modern mathematical physics. vol. 1. Functional analysis*. Academic, 1980.
- [38] H. A. Stone. A simple derivation of the time-dependent convective-diffusion equation for surfactant transport along a deforming interface. *Physics of Fluids A: Fluid Dynamics*, 2(1):111–112, jan 1990. doi: 10.1063/1.857686. URL <http://dx.doi.org/10.1063/1.857686>.
- [39] J.-J. Xu and H.-K. Zhao. An eulerian formulation for solving partial differential equations along a moving interface. *Journal of Scientific Computing*, 19(1/3): 573–594, 2003. doi: 10.1023/a:1025336916176. URL <http://dx.doi.org/10.1023/A:1025336916176>.
- [40] K. Yosida. *Functional Analysis*. Springer Nature, 1980. doi: 10.1007/978-3-642-61859-8. URL <http://dx.doi.org/10.1007/978-3-642-61859-8>.

INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT BERLIN, 14195 BERLIN, GERMANY
E-mail address: ana.djurджеvac@fu-berlin.de

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY. CV4 7AL. UK
E-mail address: C.M.Elliott@warwick.ac.uk

INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT BERLIN, 14195 BERLIN, GERMANY
E-mail address: ralf.kornhuber@fu-berlin.de

SCHOOL OF COMPUTING, UNIVERSITY OF LEEDS, LEEDS. LS2 9JT. UK
E-mail address: T.Ranner@leeds.ac.uk